

Jpn. J. Ent., 65(2): 426. June 25, 1997

New Records of Two Bugs (Heteroptera, Pentatomomorpha) from the Ryukyu Islands, Japan

Syôiti MIYAMOTO

3-34-4, Iikura, Sawara-ku, Fukuoka, 814-01 Japan

and

Masami HAYASHI

Department of Biology, Faculty of Education,
Saitama University, Urawa, 338 Japan

Key words: *Capyella horni*; *Spermatodes variolosus*; Pentatomomorpha;
new records; Japan.

Capyella horni BREDDIN, 1907 [Berytidae]

This stilt bug, belonging to the Berytinae, has hitherto been known to occur in China (Fujian and Guangdong) and Sri Lanka, and it is newly collected from Iriomote of the Ryukyus.

Specimen examined. 1 ♀, Mombanare, Ôtomi, Iriomote Is., 25. VI. 1996, M. HAYASHI *et al.*

Spermatodes variolosus (WALKER, 1867) [Pentatomidae]

This pentatomid has been known from China, Myanmar (Burma), etc., and is recorded for the first time from Miyako and Yonaguni of the Sakishima group, the Ryukyus, Japan. This bug resembles *Sepontiella aenea* (DISTANT) in dorsal habitus.

Specimens examined. 1 ♂, Cape Higashi-hennazaki, Gusukube, Miyako Is., 11. X. 1993, M. HAYASHI *et al.*; 2 ♀, Sonai, Yonaguni Is., 29. VI. 1996, M. HAYASHI *et al.*

(Received January 29, 1997; Accepted April 17, 1997)